

Yang-Lee zeros of the one-dimensional Q -state Potts model

Seung-Yeon Kim*

*School of Computational Sciences, Korea Institute for Advanced Study,
207-43 Cheongryangri-dong, Dongdaemun-gu, Seoul 130-722, Korea*

Abstract

The distributions of the Yang-Lee zeros of the ferromagnetic and antiferromagnetic Q -state Potts models in one dimension are studied for arbitrary Q and temperature. The Yang-Lee zeros of the Potts antiferromagnet have been fully investigated for the first time. The distributions of the Yang-Lee zeros show a variety of different shapes. Some of the Yang-Lee zeros lie on the positive real axis even for $T > 0$. For the ferromagnetic model this happens only for $Q < 1$, while there exist some zeros of the antiferromagnetic model on the positive real axis both for $Q < 1$ and for $Q > 1$.

PACS numbers: 05.50.+q; 64.60.Cn; 75.10.Hk

Keywords: Potts model; Partition function zeros; Ferromagnetic Yang-Lee zeros; Antiferromagnetic Yang-Lee zeros

* Electronic address: sykim@kias.re.kr

I. INTRODUCTION

The Q -state Potts model [1, 2, 3, 4, 5] is a generalization of the Ising ($Q = 2$) model. The Q -state Potts model exhibits a rich variety of critical behavior and is very fertile ground for the analytical and numerical investigations of first- and second-order phase transitions. The Potts model is also related to other outstanding problems in physics and mathematics. Fortuin and Kasteleyn [6, 7] have shown that the Q -state Potts model in the limit $Q \rightarrow 1$ defines the problem of bond percolation. They [7] also showed that the problem of resistor network is related to a $Q = 0$ limit of the partition function of the Potts model. In addition, the zero-state Potts model describes the statistics of treelike percolation [8], and is equivalent to the undirected Abelian sandpile model [9]. The $Q = \frac{1}{2}$ state Potts model has a connection to a dilute spin glass [10]. The Q -state Potts model with $0 \leq Q \leq 1$ describes transitions in the gelation and vulcanization processes of branched polymers [11]. The partition function of the Potts model is also known as the Tutte dichromatic polynomial [12] or the Whitney rank function [13] in graph theory and combinatorics of mathematics.

By introducing the concept of the zeros of the partition function in the *complex* magnetic-field plane, Yang and Lee [14] proposed a mechanism for the occurrence of phase transitions in the thermodynamic limit and yielded a new insight into the unsolved problem of the ferromagnetic Ising model in an arbitrary nonzero external magnetic field. It has been known [14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24] that the distribution of the zeros of a model determines its critical behavior. Lee and Yang [15] also formulated the celebrated circle theorem which states that the partition function zeros of the Ising ferromagnet lie on the unit circle in the complex magnetic-field ($x = e^{\beta H}$) plane. However, for the ferromagnetic Q -state Potts model with $Q > 2$ the Yang-Lee zeros in the complex x plane lie close to, but not on, the unit circle with the two exceptions of the critical point $x = 1$ ($H = 0$) itself and the zeros in the limit $T = 0$ [19, 25, 26, 27, 28]. It has been shown [27, 28] that the distributions of the ferromagnetic Yang-Lee zeros for $Q > 1$ have similar properties independent of dimension.

Recently, the exact results on the Yang-Lee zeros of the ferromagnetic Potts model have been found using the one-dimensional model. Mittag and Stephen [29] studied the Yang-Lee zeros of the three-state Potts ferromagnet in one dimension. Glumac and Uzelac [30] found the eigenvalues of the transfer matrix of the one-dimensional Potts model for general Q .

In particular, they have shown that the ferromagnetic Yang-Lee zeros can lie on the real axis for $Q < 1$. However, the properties of the Yang-Lee zeros of the *antiferromagnetic* Potts model have never been known except for the one-dimensional Ising antiferromagnet [31]. In this paper, we study the ferromagnetic and antiferromagnetic Yang-Lee zeros of the one-dimensional Q -state Potts model.

II. PARTITION FUNCTION

The Q -state Potts model in an external magnetic field H_q on a lattice G with N_s sites and N_b bonds is defined by the Hamiltonian

$$\mathcal{H}_Q = -J \sum_{\langle i,j \rangle} \delta(\sigma_i, \sigma_j) - H_q \sum_k \delta(\sigma_k, q), \quad (1)$$

where J is the coupling constant (ferromagnetic model for $J > 0$ and antiferromagnetic model for $J < 0$), $\langle i, j \rangle$ indicates a sum over nearest-neighbor pairs, $\sigma_i = 1, 2, \dots, Q$, and q is a fixed integer between 1 and Q . The partition function of the model is

$$Z_Q = \sum_{\{\sigma_n\}} e^{-\beta \mathcal{H}_Q}, \quad (2)$$

where $\{\sigma_n\}$ denotes a sum over Q^{N_s} possible spin configurations and $\beta = (k_B T)^{-1}$. The partition function can be written as

$$Z(a, x, Q) = \sum_{E=0}^{N_b} \sum_{M=0}^{N_s} \Omega_Q(E, M) a^E x^M, \quad (3)$$

where $a = y^{-1} = e^{\beta J}$, $x = e^{\beta H_q}$, E and M are positive integers $0 \leq E \leq N_b$ and $0 \leq M \leq N_s$, respectively, and $\Omega_Q(E, M)$ is the number of states with fixed E and fixed M . The states with $E = 0$ ($E = N_b$) correspond to the antiferromagnetic (ferromagnetic) ground states. For ferromagnetic interaction $J > 0$ the physical interval is $1 \leq a \leq \infty$ ($\infty \geq T \geq 0$), whereas for antiferromagnetic interaction $J < 0$ the physical interval $0 \leq a \leq 1$ ($0 \leq T \leq \infty$). The parameter Q enters the Potts model as an integer. However, the study of the Q -state Potts model has been extended to continuous Q due to the Fortuin-Kasteleyn representation of the partition function [6, 7] and its extension [32].

For the one-dimensional Potts model in an external field the eigenvalues of the transfer matrix were found by Glumac and Uzelac [30]. The eigenvalues are $\lambda_{\pm} = (A \pm iB)/2$, where

$A = a(1+x) + Q - 2$ and $B = -i\sqrt{[a(1-x) + Q - 2]^2 + 4(Q-1)x}$, and $\lambda_0 = a - 1$ which is $(Q-2)$ -fold degenerate. The partition function of the one-dimensional model ($N = N_s = N_b$) is given by

$$Z_N(a, x, Q) = \lambda_+^N + \lambda_-^N + (Q-2)\lambda_0^N. \quad (4)$$

III. PARTITION FUNCTION ZEROS

When λ_+ and λ_- are two dominant eigenvalues, the partition function becomes

$$Z_N \simeq \lambda_+^N + \lambda_-^N \quad (5)$$

for large N . If we define $A = 2C \cos \psi$ and $B = 2C \sin \psi$, where $C = \sqrt{(a-1)(a+Q-1)x}$, then $\lambda_{\pm} = C \exp(\pm i\psi)$, and the partition function is

$$Z_N = 2C^N \cos N\psi. \quad (6)$$

The zeros of the partition function are then given by

$$\psi = \psi_k = \frac{2k+1}{2N}\pi, \quad k = 0, 1, 2, \dots, N-1. \quad (7)$$

In the thermodynamic limit the locus of the partition function zeros is determined by the solution of

$$A = 2C \cos \psi, \quad (8)$$

where $0 \leq \psi \leq \pi$. In the special case $Q = 2$ the contribution by the eigenvalue λ_0 disappears from the partition function, Eq. (4), and the equation (8) determines all the locus even for finite systems. From Eq. (8) the locus of the Yang-Lee zeros for any Q is obtained to be

$$x_1(\psi) = \frac{1}{a^2} \left[\sqrt{f_1} \cos \psi + i\sqrt{f_2} \right]^2, \quad (9)$$

where $f_1 = (a-1)(a+Q-1)$ and $f_2 = f_1 \sin^2 \psi + Q - 1$. The edge zeros of $x_1(\psi)$ are given by

$$x_{\pm} = \frac{1}{a^2} \left[\sqrt{(a-1)(a+Q-1)} \pm \sqrt{1-Q} \right]^2 \quad (10)$$

from $x_1(0)$ and $x_1(\pi)$. If $f_1 > 0$ and $f_2 > 0$ or $f_1 < 0$ and $f_2 < 0$, it is easily verified that

$$|x_1(\psi)| = \sqrt{x_+ x_-} = \left| \frac{a+Q-2}{a} \right|. \quad (11)$$

From Eq. (11) we see that the zeros of $x_1(\psi)$ lie on a circle in the complex x plane. The one point of the circle $x_1(\psi)$,

$$x_1\left(\frac{\pi}{2}\right) = -\frac{a+Q-2}{a}, \quad (12)$$

always lies on the real axis. However, if $f_1 > 0$ and $f_2 < 0$ or $f_1 < 0$ and $f_2 > 0$, the zeros of $x_1(\psi)$ lie on the real axis. Recently, Ghulghazaryan et al. [33] tried to understand the properties of Eq. (9) in the antiferromagnetic ($a \leq 1$) region.

On the other hand, when λ_+ and λ_0 are two dominant eigenvalues, the partition function can be written as

$$Z_N \simeq \lambda_+^N + (Q-2)\lambda_0^N \quad (13)$$

for large N . The partition function zeros are then determined by

$$\frac{\lambda_+}{\lambda_0} = (2-Q)^{1/N} \exp(i\phi), \quad (14)$$

where

$$\phi = \phi_k = \frac{2\pi k}{N}, \quad k = 0, 1, 2, \dots, N-1. \quad (15)$$

In the thermodynamic limit the locus of the partition function zeros is determined by the solution of

$$a^2x^2 + (Q-1)x - axA + (a-1)Ae^{i\phi} - (a-1)^2e^{2i\phi} = 0, \quad (16)$$

where $0 \leq \phi \leq 2\pi$. The equation (16) also determines the locus of the zeros when λ_- and λ_0 are two dominant eigenvalues. Eq. (16) gives the second locus of the Yang-Lee zeros

$$x_2(\phi) = \frac{e^{i\phi}[a+Q-2-(a-1)e^{i\phi}]}{a+Q-1-ae^{i\phi}}, \quad (17)$$

which is the unit circle in the limit $a \rightarrow \infty$ and a type of limaçon of Pascal for $a = 0$. The two points of the locus $x_2(\phi)$,

$$x_2(0) = 1 \quad (18)$$

and

$$x_2(\pi) = -\frac{2a+Q-3}{2a+Q-1}, \quad (19)$$

can be lie on the real axis. From $|\lambda_{\pm}| = |\lambda_0|$ we also obtain

$$x_* = \frac{a-1}{a+Q-1}. \quad (20)$$

The second locus, Eq. (17), of the Yang-Lee zeros, although it is very important in understanding the Q -state Potts model, has never been considered in the literature until now.

IV. FERROMAGNETIC ($a \geq 1$) YANG-LEE ZEROS

In this section we study the Yang-Lee zeros of the ferromagnetic Q -state Potts model for $a \geq 1$.

A. $Q > 1$

In this case f_1 and f_2 are always positive and the Yang-Lee zeros $x_1(\psi)$ lie on a circle where the eigenvalues satisfy

$$\frac{|\lambda_{\pm}|}{|\lambda_0|} = \sqrt{\frac{(a+Q-1)(a+Q-2)}{a(a-1)}} > 1, \quad (21)$$

which implies that the locus $x_2(\phi)$ does not appear. For $1 < Q < 2$ the zeros lie inside the unit circle while for $Q > 2$ the zeros lie outside the unit circle, as shown in [27, 30]. In the special case $Q = 2$ we of course find that $|x_1(\psi)| = 1$, as proved by Lee and Yang [15].

The argument θ of $x_1(\psi)$ ($= |x_1(\psi)|e^{i\theta(\psi)}$) is given by

$$\cos \frac{\theta}{2} = \sqrt{\frac{(a-1)(a+Q-1)}{a(a+Q-2)}} \cos \psi. \quad (22)$$

As $a \rightarrow \infty$ ($T \rightarrow 0$), the zeros approach the unit circle. For $a = 1$ ($T = \infty$), $\cos(\theta/2) = 0$, so $\theta(\psi) = \pi$ and all the zeros lie at $1 - Q$. As $Q \rightarrow \infty$, the radius $|x_1(\psi)|$ increases without bound. The Yang-Lee edge zero for $T > 0$ is given by

$$\theta_0 = 2 \cos^{-1} \sqrt{\frac{(a-1)(a+Q-1)}{a(a+Q-2)}} > 0, \quad (23)$$

while $\theta_0 = 0$ at $T = 0$. Therefore, we conclude that no zero lie on the positive real axis for any $T > 0$.

B. $Q < 1$

At $a = 1$ ($T = \infty$), all the Yang-Lee zeros lie at the point $1 - Q$ on the positive real axis. For $1 < a \leq 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$, all the zeros lie on the positive real axis between x_- and x_+ ($0 < x_- < x_+ < 1$), as pointed out by Glumac and Uzelac [30, 33, 34]. For $a > 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$, the locus consists of the loop $x_2(\phi)$ ($\phi_* \leq \phi \leq 2\pi - \phi_*$) and the line $x_1(\psi)$ ($0 \leq \psi \leq \psi_*$) on

the positive real axis between $x_* = x_1(\psi_*)$ and $x_+ = x_1(0)$ ($0 < x_* < x_+ < 1$). The loop x_2 meets with the line x_1 at the point

$$x_* = x_2(\phi_*) = x_2(2\pi - \phi_*), \quad (24)$$

where $|\lambda_+| = |\lambda_-| = |\lambda_0|$. The loop $x_2(\phi)$ cuts the real axis at two points $x_2(\pi)$ and x_* ($> x_2(\pi)$). The sign of $x_2(\pi)$ is positive for $1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2} < a < \frac{3-Q}{2}$ and negative for $a > \frac{3-Q}{2}$. Figure 1 shows the locus of the Yang-Lee zeros for $Q = \frac{1}{2}$ and $a = \frac{6}{5}$ from which we obtain $x_2(\pi) = \frac{1}{19}$, $x_* = \frac{2}{7}$, and $x_+ = 0.8119$.

At $Q = 0$, the line $x_1(\psi)$ shrink to the point

$$x_* = x_+ = x_2(0) = 1, \quad (25)$$

and all the zeros lie on the loop $x_2(\phi)$ ($0 \leq \phi \leq 2\pi$) for $a > 1$. As $Q \rightarrow 1$, two points x_* and x_+ approach $(a-1)/a$ together, and the line $x_1(\psi)$ disappears. At $a = \infty$ ($T = 0$), the line $x_1(\psi)$ again shrink to the point $x_* = x_+ = 1$, and the loop $x_2(\phi)$ becomes the unit circle centered at the origin.

V. ANTIFERROMAGNETIC ($0 \leq a \leq 1$) YANG-LEE ZEROS

In this section we investigate the Yang-Lee zeros of the antiferromagnetic Q -state Potts model for $0 \leq a \leq 1$.

A. $Q > 1$

Because f_1 is always negative, the zeros of $x_1(\psi)$ lie on a circle if $f_2 < 0$ ($\psi_0 < \psi < \pi - \psi_0$) and on the real axis if $f_2 > 0$ ($0 \leq \psi < \psi_0$ or $\pi - \psi_0 < \psi \leq \pi$). For $a < 1 - \frac{Q}{2}$, the locus consists of the line $x_1(\psi)$ ($0 \leq \psi \leq \psi_0$ and $\pi - \psi_0 \leq \psi \leq \psi_*$) on the negative real axis between $x_+ = x_1(0)$ and $x_* = x_1(\psi_*)$ ($x_+ < x_* < 0$), the circle $x_1(\psi)$ ($\psi_0 \leq \psi \leq \pi - \psi_0$) with the radius $|x_1(\psi)|$, and the loop $x_2(\phi)$ ($0 \leq \phi \leq \phi_*$ and $2\pi - \phi_* \leq \phi \leq 2\pi$), inside the circle x_1 . The loop x_2 again meets with the line x_1 at the point

$$x_* = x_2(\phi_*) = x_2(2\pi - \phi_*), \quad (26)$$

where $|\lambda_+| = |\lambda_-| = |\lambda_0|$. The circle cuts the real axis at two points x_1^0 and $x_1(\frac{\pi}{2})$ ($= -x_1^0 > 1$), where the point x_1^0 is defined by

$$x_1^0 = x_1(\psi_0) = x_1(\pi - \psi_0) = \frac{a + Q - 2}{a}. \quad (27)$$

Similarly, the loop also cuts the real axis at two points x_* (< 0) and $x_2(0)$ ($= 1$). At $a = 0$ ($T = 0$), the circle $x_1(\psi)$ disappears for $1 < Q < 2$, and the locus consists of the line $x_1(\psi)$ on the real axis between $-\infty$ and x_* (< 0) and the loop $x_2(\phi)$. For example, figure 2 shows the locus for $Q = \frac{11}{10}$ and $a = \frac{1}{10}$. In this case we obtain $x_+ = -54.83$, $x_1^0 = -8$, $x_* = -\frac{9}{2}$, the radius $|x_1| = 8$ for the circle, $\psi_0 = 48.19^\circ$, $\psi_* = 134.07^\circ$, and $\phi_* = 46.02^\circ$.

At $a = 1 - \frac{Q}{2}$, two points x_* and x_1^0 on the real axis meet at

$$x_* = x_1^0 = -1, \quad (28)$$

other two points $x_1(\frac{\pi}{2})$ and $x_2(0)$ on the real axis also meet at

$$x_1\left(\frac{\pi}{2}\right) = x_2(0) = 1, \quad (29)$$

and the circle $x_1(\psi)$ and the loop $x_2(\phi)$ become the identical locus as the unit circle. On this unit circle, three eigenvalues have the same magnitude

$$|\lambda_+| = |\lambda_-| = |\lambda_0| = \frac{Q}{2}. \quad (30)$$

Therefore, the locus consists of the line $x_1(\psi)$ on the real axis between x_+ and x_* and the circle.

In the region $1 - \frac{Q}{2} < a < 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$, the circle $x_1(\psi)$ disappears, and the line $x_1(\psi)$ ($0 \leq \psi \leq \psi_*$) on the negative real axis between x_+ and x_* ($x_+ < x_* < 0$) again meets with the loop $x_2(\phi)$ ($\phi_* \leq \phi \leq 2\pi - \phi_*$) at the point x_* . The loop cuts the real axis at two points x_* and $x_2(\pi)$. The sign of $x_2(\pi)$ is positive ($0 < x_2(\pi) < 1$) for $1 - \frac{Q}{2} < a < \frac{3-Q}{2}$ and negative for $\frac{3-Q}{2} < a < 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$. Figure 3 shows the locus of the Yang-Lee zeros for $Q = 3$ and $a = \frac{1}{10}$ which give $x_2(\pi) = -\frac{1}{11}$, $x_* = -\frac{3}{7}$, and $x_+ = -777.84$. At $a = 0$ ($T = 0$), the locus still consists of the line $x_1(\psi)$ on the real axis between $-\infty$ and x_* (< 0) and the loop $x_2(\phi)$ for $2 < Q < 4$. At $T = 0$, the sign of $x_2(\pi)$ is positive ($0 \leq x_2(\pi) < 1$) for $2 < Q \leq 3$ and negative for $3 < Q < 4$. As $Q \rightarrow 1$, two points x_+ and x_* approach $(a - 1)/a$ together, and the line $x_1(\psi)$ disappears.

At $a = 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$, $x_* = x_-$, and the loop $x_2(\phi)$ shrinks to the point x_* . For $a > 1 - \frac{Q}{2} + \frac{\sqrt{Q}}{2}$, the loop disappears, and the only locus is the line $x_1(\psi)$ on the negative real axis between

x_+ and x_- ($x_+ < x_- < 0$). As $Q \rightarrow \infty$, $x_{\pm} \rightarrow -\infty$ for $a < 1$. At $a = 1$ ($T = \infty$), two edge zeros x_+ and x_- meet, and the line $x_1(\psi)$ shrinks to the point

$$x_+ = x_- = 1 - Q \quad (31)$$

for $Q > 1$. In the special case $Q = 2$, $f_1 < 0$ and $f_2 > 0$, and the line $x_1(\psi)$ is the only locus of the Yang-Lee zeros. Therefore, all the Yang-Lee zeros of the antiferromagnetic Ising model lie on the negative real axis between x_+ and x_- ($> x_+$) for $a < 1$, as shown by Yang [31]. At $a = 0$ ($T = 0$), all the zeros lie on the negative real axis between two edge zeros,

$$x_a = -\infty \quad (32)$$

and

$$x_b = \frac{(Q-2)^2}{4(1-Q)} \quad (x_b \leq 0), \quad (33)$$

for $Q = 2$ and $Q \geq 4$.

B. $Q < 1$

For $0 \leq a < 1 - \frac{Q}{2} - \frac{\sqrt{Q}}{2}$, the locus consists of the loop $x_2(\phi)$ ($0 \leq \phi \leq \phi_*$ and $2\pi - \phi_* \leq \phi \leq 2\pi$) and the line $x_1(\psi)$ ($0 \leq \psi \leq \psi_*$) on the positive real axis between $x_* = x_1(\psi_*)$ and $x_+ = x_1(0)$ ($1 < x_* < x_+$). The loop x_2 cuts the real axis at two points $x_2(0)$ ($= 1$) and x_* (> 1). The loop again meets with the line x_1 at the point

$$x_* = x_2(\phi_*) = x_2(2\pi - \phi_*), \quad (34)$$

where $|\lambda_+| = |\lambda_-| = |\lambda_0|$. At $Q = 0$, two points $x_2(0)$ and x_* meet, the loop $x_2(\phi)$ shrinks to the point $x_2(0) = x_*$, and all the Yang-Lee zeros lie on the positive real axis between x_- ($= x_* = 1$) and x_+ ($\geq x_-$) for $a \leq 1$.

At $a = 1 - \frac{Q}{2} - \frac{\sqrt{Q}}{2}$, $x_* = x_+$, and the line $x_1(\psi)$ shrinks to the point x_* . In the region $1 - \frac{Q}{2} - \frac{\sqrt{Q}}{2} < a < 1 - \frac{Q}{2}$, the line disappears, and the locations of Yang-Lee zeros are completely determined by the locus $x_2(\phi)$ ($0 \leq \phi \leq 2\pi$) which cuts the real axis at two points $x_2(0)$ ($= 1$) and $x_2(\pi)$ (> 1). The shape of the locus $x_2(\phi)$ is changed from a waterdrop-like shape for $1 - \frac{Q}{2} - \frac{\sqrt{Q}}{2} < a < 1 - Q$ through a circle with center $\frac{1}{1-Q}$ and radius $\frac{Q}{1-Q}$ for $a = 1 - Q$ to a crescent-like shape for $1 - Q < a < 1 - \frac{Q}{2}$. Figure 4 shows the locus of the Yang-Lee zeros for $Q = \frac{9}{10}$ and $a = \frac{2}{5}$ which give $x_2(\pi) = \frac{13}{7}$.

At $a = 1 - \frac{Q}{2}$, the loci $x_1(\psi)$ and $x_2(\phi)$ become identical to be the unit circle on which three eigenvalues again have the same magnitude

$$|\lambda_+| = |\lambda_-| = |\lambda_0| = \frac{Q}{2}. \quad (35)$$

For $a > 1 - \frac{Q}{2}$, all the Yang-Lee zeros lie on the circle $x_1(\psi)$ ($0 \leq \psi \leq \pi$) which cuts the positive real axis at the point $x_1(\frac{\pi}{2})$ ($0 < x_1(\frac{\pi}{2}) < 1$). At $a = 1$ ($T = \infty$), this circle shrinks to the point $x_1(\psi) = 1 - Q$ for $Q < 1$.

VI. ONE-STATE (Q=1) POTTS MODEL

At $Q = 1$, the partition function becomes

$$Z_N(a, x, Q = 1) = (ax)^N. \quad (36)$$

Therefore, all the Yang-Lee zeros lie on the point $x = 0$. On the other hand, in the limit $Q \rightarrow 1$, the line $x_1(\psi)$ disappears, and the circle $x_1(\psi)$ is the uniform distribution of the zeros because the circle approaches

$$x_1(\psi) = \left| \frac{a-1}{a} \right| e^{2i\psi}. \quad (37)$$

The loop $x_2(\phi)$ also approaches

$$x_2(\phi) = \frac{a-1}{a} e^{i\phi}, \quad (38)$$

which is the same as the circle $x_1(\psi)$.

VII. CONCLUSION

We have studied the interesting properties of the ferromagnetic and antiferromagnetic Yang-Lee zeros of the one-dimensional Q -state Potts model for arbitrary Q and temperature. It has been shown that the distributions of the Yang-Lee zeros have a variety of different shapes. In particular, the antiferromagnetic Yang-Lee zeros of the Potts model have been fully investigated for the first time. One of the most interesting results is that some of the Yang-Lee zeros lie on the positive real axis even for $T > 0$. For the ferromagnetic Yang-Lee zeros this happens only for $Q < 1$, while there exist some Yang-Lee zeros of the antiferromagnetic Potts model on the positive real axis both for $Q < 1$ and for $Q > 1$.

The results obtained from the one-dimensional Potts model may be considered as a road to the full understanding of the Yang-Lee zeros in higher dimensions in that for $Q > 1$ the distributions of the ferromagnetic Yang-Lee zeros have similar properties independent of dimension [27, 28].

Acknowledgments

The author is grateful to Prof. R.J. Creswick for very useful discussions.

-
- [1] R.B. Potts, Proc. Cambridge Philos. Soc. 48 (1952) 106.
 - [2] F.Y. Wu, Rev. Mod. Phys. 54 (1982) 235.
 - [3] F.Y. Wu, J. Appl. Phys. 55 (1984) 2421.
 - [4] R.J. Baxter, Exactly Solved Models in Statistical Mechanics, Academic Press, London, 1982.
 - [5] P.P. Martin, Potts Models and Related Problems in Statistical Mechanics, World Scientific, Singapore, 1991.
 - [6] P.W. Kasteleyn, C.M. Fortuin, J. Phys. Soc. Jpn. Suppl. 26 (1969) 11.
 - [7] C.M. Fortuin, P.W. Kasteleyn, Physica 57 (1972) 536.
 - [8] M.J. Stephen, Phys. Lett. A 56 (1976) 149.
 - [9] S.N. Majumdar, D. Dhar, Physica A 185 (1992) 129.
 - [10] A. Aharony, J. Phys. C 11 (1978) L457.
 - [11] T.C. Lubensky, J. Isaacson, Phys. Rev. Lett. 41 (1978) 829.
 - [12] W.T. Tutte, Proc. Cambridge Philos. Soc. 43 (1947) 26.
 - [13] H. Whitney, Ann. Math. 33 (1932) 688.
 - [14] C.N. Yang, T.D. Lee, Phys. Rev. 87 (1952) 404.
 - [15] T.D. Lee, C.N. Yang, Phys. Rev. 87 (1952) 410.
 - [16] C. Itzykson, R.B. Pearson, J.B. Zuber, Nucl. Phys. B 220 (1983) 415.
 - [17] R.J. Creswick, S.-Y. Kim, Phys. Rev. E 56 (1997) 2418.
 - [18] R.J. Creswick, S.-Y. Kim, in: D.P. Landau, K.K. Mon, H.-B. Schüttler (Eds.), Computer Simulation Studies in Condensed-Matter Physics, Vol. 10, Springer, Berlin, 1998, p. 224.
 - [19] R.J. Creswick, S.-Y. Kim, Comput. Phys. Commun. 121 (1999) 26.
 - [20] W. Janke, R. Kenna, J. Stat. Phys. 102 (2001) 1211.
 - [21] W. Janke, R. Kenna, Comput. Phys. Commun. 147 (2002) 443.
 - [22] W. Janke, R. Kenna, Nucl. Phys. B (Proc. Suppl.) 106 (2002) 905.
 - [23] N.A. Alves, J.P.N. Ferrite, U.H.E. Hansmann, Phys. Rev. E 65 (2002) 036110.
 - [24] R.A. Kromhout, Physica B 301 (2001) 389.
 - [25] S.-Y. Kim, R.J. Creswick, Phys. Rev. Lett. 81 (1998) 2000.
 - [26] S.-Y. Kim, R.J. Creswick, Phys. Rev. Lett. 82 (1999) 3924.
 - [27] S.-Y. Kim, R.J. Creswick, Physica A 281 (2000) 252.

- [28] S.-Y. Kim, Nucl. Phys. B 637 (2002) 409.
- [29] L. Mittag, M.J. Stephen, J. Stat. Phys. 35 (1984) 303.
- [30] Z. Glumac, K. Uzelac, J. Phys. A 27 (1994) 7709.
- [31] C.N. Yang, Special Problems of Statistical Mechanics (Lectures at the University of Washington), 1952.
- [32] H.W.J. Blöte, M.P. Nightingale, Physica A 112 (1982) 405.
- [33] R.G. Ghulghazaryan, N.S. Ananikan, P.M.A. Sloot, Phys. Rev. E 66 (2002) 046110.
- [34] J.L. Monroe, J. Phys. A 29 (1996) 5421.




FIG. 1: The locus of the Yang-Lee zeros in the complex $x = e^{\beta H_q}$ plane for $Q = \frac{1}{2}$ and $a = e^{\beta J} = \frac{6}{5}$. For comparison, the zeros for a finite-size system ($N = 100$) are also shown (open circles).




FIG. 2: The locus of the Yang-Lee zeros for $Q = \frac{11}{10}$ and $a = \frac{1}{10}$.




FIG. 3: The locus of the Yang-Lee zeros for $Q = 3$ and $a = \frac{1}{10}$. The zeros on the real axis lie between $x_* = -\frac{3}{7}$ and $x_+ = -777.84$. Most of them are omitted in the figure.




FIG. 4: The locus of the Yang-Lee zeros for $Q = \frac{9}{10}$ and $a = \frac{2}{5}$.